



Characterization of soliton compounds in a passively mode-locked high power fiber laser

Submitted by Emmanuel Lemoine on Wed, 10/29/2014 - 11:43

Titre	Characterization of soliton compounds in a passively mode-locked high power fiber laser
Type de publication	Article de revue
Auteur	Amrani, Foued [1], Salhi, Mohamed [2], Leblond, Hervé [3], Sanchez, François [4]
Editeur	Elsevier
Type	Article scientifique dans une revue à comité de lecture
Année	2010
Langue	Anglais
Date	2010/12/15
Numéro	24
Pagination	5224 - 5230
Volume	283
Titre de la revue	Optics Communications
ISSN	0030-4018
Résumé en anglais	<p>We characterize soliton complexes in a high power double-clad erbium-doped fiber laser passively mode-locked through nonlinear polarization evolution. Such complexes involve some hundreds of solitons and form self-organized or disorganized patterns analogous to the states of the matter. Experimentally these soliton compounds are characterized through the autocorrelation trace, the optical spectrum and the oscilloscope trace which is limited due to its finite bandwidth. We perform here a reconstruction of the experimental results thus allowing us to identify the temporal distribution of the solitons inside the cavity. The reconstruction allows us to clarify and either to confirm or to correct the former intuitive interpretation. Especially, a soliton 'spray' is identified.</p>
URL de la notice	http://okina.univ-angers.fr/publications/ua5113 [5]
DOI	10.1016/j.optcom.2010.08.003 [6]
Lien vers le document	http://dx.doi.org/10.1016/j.optcom.2010.08.003 [6]

Liens

- [1] [http://okina.univ-angers.fr/publications?f\[author\]=8557](http://okina.univ-angers.fr/publications?f[author]=8557)
- [2] <http://okina.univ-angers.fr/m.salhi/publications>
- [3] <http://okina.univ-angers.fr/herve.leblond/publications>
- [4] <http://okina.univ-angers.fr/francois.sanchez/publications>
- [5] <http://okina.univ-angers.fr/publications/ua5113>
- [6] <http://dx.doi.org/10.1016/j.optcom.2010.08.003>

